

Prevalence of Oral Manifestations and Associated Factors in the HIV Infection at a Reference Hospital from the Health's Unified System, Brazil – an update

Solange M. Silva^{1,*}, Rejane F. Ribeiro-Rotta¹ and Marco Tulio A. García-Zapata²

¹Avenida Universitária com 1ª Avenida, s/n. Faculdade de Odontologia – Praça Universitária s/n. Setor Universitário, Goiânia, Zip Code 74605 220, Brazil

²Rua 235 s/n. Setor Universitário, Instituto de Patologia Tropical e Saúde Pública – IPTSP, Zip Code 74605, Brazil

Abstract: A severe control by the Brazilian Ministry of Health on the HIV infection suggests that all the opportunistic infections related to this disease are in a low frequency. This study aimed to characterize the oral lesions profile in patients with HIV/AIDS who presented to the public health system in Goiânia, Brazil. An oral clinical examination was developed based on the EC-Clearinghouse on oral problems related to HIV infection. A questionnaire was completed, medical data were collected and photos were taken. A total of 201 patients were divided into two groups: Group I (92 HIV-positive patients with CD4+ count above 200 cells/mm³ and absence of symptoms of AIDS) and Group II (109 HIV-positive patients with CD4+ count below 200 cells/mm³ and with signs and symptoms of AIDS). The total prevalence for oral manifestations ranged from 19.60% (Group I) to 33.00% (Group II). Periodontitis associated with HIV infection occurred in 5.43% (Group I) and 11.00% (Group II) of patients. Linear gingival erythema occurred in 4.34% (Group I) and 7.34% (Group II) of patients. Erythematous candidiasis affected 4.34% (Group I) and 2.75% (Group II) of patients. Pseudomembranous candidiasis affected 1.83% of Group I patients. A significant association was observed between the occurrence of oral lesions related to HIV infection and age ($p < 0.05$; OR 2.82; 95% CI (1.31-6.05)) and between HIV-related oral lesions and prosthesis wearing ($p < 0.05$; OR 3.6; 95% CI (1.2-11.4)) in Group I patients. In Group II patients, an association was observed between the presence of caries and HIV-related oral lesions ($p < 0.005$; OR 2.4; 95% CI (1.1-5.4)). Dry mouth was the most observed complaint related to antiretroviral therapy (7.80% in Group I and 11.00% in Group II). In conclusion, the Group II presented the highest prevalence of lesions, and periodontitis was the most common oral lesion. Age and prosthesis wearing were factors associated with the development of these lesions among Group I patients, while the presence of dental caries was associated with oral lesions in Group II patients.

Keywords: Oral manifestations, mouth disease, AIDS-related opportunistic infections, HIV.

INTRODUCTION

Oral manifestations of HIV infection represent a major proportion of the opportunistic infections that afflict these patients and affect their quality of life [1]. These manifestations are not only important in the early diagnosis of HIV infection but also in monitoring a patient's progression to AIDS. The majority of cases can be diagnosed by their clinical features alone [2].

More than 40 types of oral lesions have been associated with a greater extent of HIV infection. The aetiological agents for HIV-associated oral lesions include fungal infections (candidiasis, linear gingival erythema and deep mycoses), bacterial infections (necrotizing gingivitis, necrotizing periodontitis, bacillary angiomatosis and tuberculosis), viral infections (herpes simplex, hairy leukoplakia, warts, herpes zoster and cytomegalovirus ulcers), neoplasms (Kaposi's sarcoma and oral lymphoma) and other causes (nonspecific ulcers, xerostomia, parotid swelling and intraoral

pigmentation). It is essential to emphasize that these lesions may also be present in the general population. However, in patients with HIV/AIDS, these lesions have an aggressive, persistent and recurrent behavior [3].

Because there are increasing numbers of patients living with the infection, the number of HIV-positive patients who may seek dental care is increasing. Oral examinations may assist in the early recognition of the disease and aid in the comprehensive evaluation of HIV-infected patients [4].

Strategies for training health-care professionals in the diagnosis of the oral manifestations of HIV are important and necessary to prevent pain, discomfort, malnutrition, weight loss, dehydration, dry mouth, non-adherence to antiretroviral therapy and, sometimes, the eventual death of these patients [5].

The reference centers in diagnosis of oral lesions and preventive measures for these lesions, like oral hygiene instructions and an available access to oral treatment should be part of HIV-patients management developed for a multiprofessional team [6,7].

*Address correspondence to this author at the Avenida Universitária com 1ª Avenida, s/n. Faculdade de Odontologia – Praça Universitária s/n. Setor Universitário, Goiânia, Zip Code 74605 220, Brazil; Tel: 55 (62) 84246648; Fax: 55 (62) 3209 6067; E-mail: alecrims@gmail.com

It is becoming increasingly challenging for clinicians to accurately diagnose new lesions, due to the vast array of manifestations of AIDS in the neck and head region and their atypical presentations [8, 9].

Even with the Highly Active Antiretroviral Therapy (HAART) advent, HIV positive patients continue to experience high levels of oral manifestations –HIV and still have oral health needs that impact the general health [10].

There are few data on the oral manifestations of HIV/AIDS among patients in Brazil, a country considered as a model for addressing this infection. Therefore, this study was designed to characterize the profile of oral health among patients located in an important unit referenced by the Unified Health System (SUS). This investigation also intends to emphasize the value in monitoring these oral manifestations to assess the immune status of patients with HIV/AIDS and identifying treatment failures.

MATERIALS AND METHODS

Patients

The study sample consisted of 201 consecutive patients of both sexes, aged 18 years and older, who presented to the public clinic of the Hospital of Clinics of the Federal University of Goiás (HC/UFG) between June 2011 and November 2012. The study sample size was calculated based on the prevalence of oral manifestations described in the literature (22.00% to 78.00%) and with an estimated margin of error of 5.00%. The total number of registered patients at the Clinic of Infectious Diseases, HC/UFG, and the daily patient volume were also considered. This hospital is recognized for its care of individuals with HIV/AIDS from all over the state of Goiás.

The classification system for HIV infection among adolescents and adults has been revised to include the CD4+ T-lymphocyte count as a marker for HIV-related immunosuppression. As the number of CD4+ T-lymphocytes decreases, the risk and severity of opportunistic illnesses increase [11]. Because of these observations, measures of CD4+ T-lymphocytes are commonly used to guide clinical and therapeutic management of HIV-infected persons in Brazil [12]. The study participants were divided into two groups to characterize a less severe phase at the infection (the Group I, which consisted of 92 HIV-positive patients with CD4+ cell counts above 200 cells/mm³, and an absence of sign and symptoms of AIDS), and a more

severe phase (Group II, which consisted of 109 HIV-positive patients with signs and symptoms of the terminal stage of the disease and CD4+ cell count below 200 cells/mm³ and a current or previous opportunistic disease that characterizes AIDS) [12].

The study protocol was approved by the Ethics Committee of HC/UFG (CEPMHA, protocol number 051/2011) and all participants assigned a Written Informed Consent (WIC). Patients with 18 years old or more and with a preserved cognitive health were included.

Oral Clinical Examination, Data from Medical Records and Research-Specific Questionnaire

To assess previously developed research tools (dental examinations, data collection from medical records and research-specific questionnaires), a pilot study was conducted with 72 patients during the period of June through November of 2011. Patients underwent a clinical examination in a private room in a regular chair under artificial light using a mouth mirror, a dental explorer, a periodontal probe and gauze. Only one dentist conducted the exams. It was used a presumptive diagnosis based on the criteria provided by the EC Clearinghouse on Oral Problems related to HIV infection [13]. Lesions were photographed with a professional camera. The presence or absence of any dental lesions, including the occurrence of dental caries, was recorded.

Diagnostic Conclusion

The diagnosis of lesions of the oral mucosa (except dental caries) was confirmed by consensus with the team of infectious disease expertise and confirmed through histopathological examinations when necessary. Xerostomia was considered by the viscosity of saliva and the patient's report. All patients answered a structured record of research, addressing shareholders demographic data, medical and laboratory data (recent viral load, CD4 and CD8 counts, registration AIDS defining illnesses, diagnosis time and route of HIV transmission) collected from medical records. It was registered the presence of alcohol and smoking habits, the presence of dental caries, oral hygiene habits, the use of dental prostheses, access to regular dental treatment and reporting of complaints due to use of antiretroviral therapy.

Statistical Analysis

Data were tabulated in Microsoft Excel 2007 (Microsoft ®, Brazil). Statistical analyses were

performed using SPSS for Windows (version 15.0, USA). Descriptive statistical analyses included the frequency of oral manifestations associated with HIV infection and the calculation of mean frequencies and standard deviations for any oral lesion detected. The Fisher exact test was used to compare the prevalence of oral lesions between the two groups. A univariate logistic regression analysis was used to calculate the odds ratio (OR) and its confidence interval (CI) to estimate the strength of association between the presence of HIV-related oral lesions and various traits including gender, age, race, marital status, degree of education, monthly income, use of antiretroviral regimens, CD4+ and CD8+ cell counts, smoking habits, alcohol use, dental hygiene, dental prosthesis use, opportunity for regular dental treatment and the presence of caries. A significance level of 5.00% was considered.

RESULTS

Patient Characteristics

This study enrolled 201 patients with HIV/AIDS who were registered at the Clinic of Infectious Diseases at The Hospital of Clinics in The Federal University of Goiás. The mean age was 34.38 ± 10.96 years for Group I and 40.91 ± 10.66 for Group II. The study subjects were predominantly male in both groups (75.00% in Group I and 61.50% in Group II). More than 70.00% of the subjects in both groups stated that they were single, and the route of HIV transmission was most frequently due to heterosexual relations among Group II patients (62.40%) and homosexual relations among Group I patients (51.10%). Additional characteristics of the study population are shown in Table 1.

Table 1: General Characteristics of the Population HIV / AIDS of Patients Regulated by SUS, Goiás, in the Period of 2011-2012

	Group I n= 92		Group II n= 109		Total n=201	
	n	%	n	%	N	%
Sex						
Male	69	75,0	67	61,5	136	67,66
Female	23	25,0	42	38,5	65	32,33
Race						
White	35	38,0	42	38,5	77	38,30
Black	18	19,6	27	24,8	45	22,38
Parda	39	42,4	40	36,7	79	39,30
Marital Status						
Married	23	25,0	32	29,4	55	27,36
Single	69	75,0	77	70,6	146	72,63
Schooling						
Fundamental	20	21,7	48	44,0	68	33,83
Medium	41	44,6	43	39,4	84	41,79
Superior	31	33,7	18	16,5	49	24,37
DiseaseDuration						
< 1 year	40	43,5	20	18,3	60	29,85
1to 5 years	39	42,4	60	55,0	99	49,25
Above5years	13	14,1	29	26,6	42	20,89
ART	24	26,1	101	92,7	125	62,18
Tobaco	20	21,7	26	24,1	46	22,88
Alcoholism	43	46,7	37	33,9	80	39,80
HIVTransmission						
Ignored	5	5,4	9	8,3	14	6,96
Homossexual	47	51,1	28	25,7	75	37,31
Bissexual	12	13,0	4	3,7	16	7,96
Heterossexual	27	29,3	68	62,4	95	47,26



Figure 1: HIV – related periodontitis (A) and Classic Periodontitis (B).

Oral Manifestations

The prevalence of oral manifestations associated with HIV infection was 19.60% and 33.00% for Groups I and II, respectively. The prevalence of any oral lesion, including those not associated with HIV infection, was 31.52% in Group I and 45.87% ($p < 0.05$). In the Group I, two patients (2.17%) had more than one oral lesion, while in the Group II, nine subjects (8.25%) had more than one lesion.

The most prevalent oral manifestation was HIV related periodontitis (5.43% in Group I and 11.00% in Group II), an aggressive inflammatory infection that leads to the rapid development of severe periodontal tissue disease (Figure 1). Necrotizing ulcerative gingivitis was uncommon (1.08% and 0.92% in Groups I and II, respectively). In Group I patients, CD4+ lymphocyte count in cases of HIV-related periodontitis varied between 281 and 741 cells/mm³, and the viral load ranged between 1487 and 185.845 viral particles/cm³. In Group II patients with HIV-related periodontitis, the CD4+ cell count ranged between 161 and 837 cells/mm³, and the viral load ranged between an undetectable level and 189 viral particle/mm³.

The second most frequent oral lesion was linear gingival erythema (Figure 2), an erythematous halo in the marginal gingiva, especially around the anterior teeth (4.34% in Group I and 7.34% in Group II).

The prevalence of erythematous candidiasis (EC) was 4.34% for Group I and 2.75% for Group II. Only Group II patients showed the pseudomembranous form

(1.83%). Angular cheilitis was present in Group I (4.34%) and Group II (1.83%) patients. The presence of erythematous candidiasis (EC) was associated with CD4+ cell counts between 443 and 741 cells/mm³ in

Group I and the viral load ranged from 10.897 to 116.231 viral particles/mm³. In Group II patients, the presence of EC was associated with CD4+ cell counts ranging from 57 to 420 cells/mm³, and the viral load ranged from an undetectable level to 231.810 viral particles/mm³. The pseudomembranous form, present only in patients of the Group II, was associated with CD4+ cell counts between 120 and 526 cells/mm³ and a viral load ranging from an undetectable level to 750 viral particles/mm³.



Figure 2: Linear Gingival Erythema.

Herpes simplex lesions were not found and Herpeszoster was observed in only 0.92% of Group I patients. Bucal hairy leukoplakia (LPB) was recorded only in Group II patients (1.08%) and was associated with a CD4+ cell count of 361 cells /mm³ and a viral load of 86.721 viral particles/mm³.



Figure 3: Carious lesions in a patient with AIDS.

Other Manifestations

Ulcers, sores, hyperpigmentation or pericoronitis were rarely observed. The dental needs of patients were important for both groups. In Group I, 38.04% of patients had at least one carious lesion, while in Group II, 41.08% of patients had caries (Figure 3). Classical periodontitis was observed in 11.95% of Group I patients and in 15.59% of Group II patients (Figure 1). Classical gingivitis affected only Group II patients (0.92%) (Table 2).

Antiretroviral Therapy and Oral Adverse Effects

Treatment with antiretroviral drugs caused unpleasant oral effects as reported by the study patients. Xerostomia was the most frequently reported adverse oral effect in both groups (7.60% in Group I and 11.00% in Group II). Antiretrovirals used by patients who reported xerostomia included zidovudine, lamivudine, efavirenz, biovir and lopinavir with ritonavir. Taste disturbance was reported by 4.30% and 5.50% of Group I and II patients, respectively. The occurrence of nausea was reported by 4.30% of patients in both groups. Nausea and taste perversion occurred with the use of the previously listed antiretroviral drugs, with the exception of lopinavir associated with ritonavir.

Association between the Occurrence of Oral Manifestations and Relevant Factors

There was no association between the presence of oral lesions and the CD4+ T-lymphocyte count, viral load, disease duration, use of antiretroviral therapy or route of HIV infection ($p > 0.05$). Age was associated with the presence of HIV-related oral manifestations ($p < 0.05$; OR 2.82; 95% CI (1.31 - 6.05)) in Group I patients

(Table 3). In this group, there was a significant association between the use of dental prostheses and the presence of oral manifestations ($p < 0.05$; OR 3.3; 95% CI (1.2 - 11.4)) (Table 4). In Group II patients, the occurrence of HIV related oral lesions was associated with the presence of dental caries ($p < 0.05$; OR 2.4; 95% CI (1.1 - 5.4)) (Table 5).

DISCUSSION

The results of this study allowed us to characterize differences in the occurrence of oral manifestations between HIV patients with a less severe infection (Group I) and patients in an advanced stage of the disease (Group II). The number of patients evaluated in this study was lower than the 605 examined patients in a study from Africa [14] and the 300 patients seen by the University Hospital of the Federal University of Rio Grande do Sul [15]. However, the sample size was above the 87 examined patients in a study from Thailand [16].

The average age was similar to Cambodian and Thai investigations [17, 18] confirming this morbidity in a productive stage of life and with an intense sexual activity.

The predominance of males in this investigation was similar to that observed in studies in Africa (58.00%), Rio Grande do Sul (58.00%) and Paraiba (58.01%) [4, 15, 19] which can be justified probably by a risk behavior adopted in this group. In contrast, an assessment of HIV/AIDS in Uganda, Africa included a higher percentage of female patients [14].

The marital status of the participants has been minimally discussed. However, the majority of Thai

Table 2: Prevalence of Oral Manifestations in HIV + / AIDS Patients, Regulated by SUS-Goiás, in the Period of 2011-2012

Oral Manifestations	Group I n= 92		Group II n= 109		p
	n	%	n	%	
HIV-RELATED ORAL MANIFESTATIONS	18	19.60	36	33.0	0,032
ANY ORAL MANIFESTATION	29	31.52	50	45.87	0,038
LESIONS WITH FUNGAL ORIGIN:					
Pseudomembranous Candidiasis	0	-	2	1.83	0,192
Erythematous Candidiasis	4	4.34	3	2.75	0,539
Angular Cheilitis	4	4.34	2	1.83	0,297
LESIONS WITH VIRAL ORIGIN:					
Herpes Zoster	0	-	1	0.92	0,357
Oral Hairy Leukoplakia	1	1.08	0	-	0,275
LESIONS WITH BACTERIAL ORIGIN:					
Linear Gingival Erythema	4	4.34	8	7.34	0,372
Necrotizing Ulcerative Gingivitis	1	1.08	1	0.92	0,904
Periodontitis HIV- Related	5	5.43	12	11.00	0,157
SALIVARY GLAND LESION:					
Ranula	1	1.08	0	-	0,275
Xerostomia	1	1.08	7	6.42	0,060
LESIONS UNRELATED TO HIV					
Classic Gingivitis	0	-	1	0.92	0,357
Classic Periodontitis	11	11.95	17	15.59	0,458
Caries	35	38.04	45	41.28	0,640
Argirose	0	-	1	0.92	0,357
OTHER LESIONS:					
Pericoronaritis	0	0	1	0.92	0,357
Úlcers	1	0,98	3	2,75	0,400
Hyperpigmentation	0	-	1	0.92	0,357

p significance level **SUS** Unified Health System.

subjects in the two evaluated groups from different regions were married (58.00% versus 26.00%) [18].

Regarding the likely route of HIV transmission, the results were similar to findings from studies conducted in a city in southern Brazil (75.00% were heterosexual) and in India (59.8% were heterosexual) [15,20]. A study at the University Hospital of Minas Gerais found one of two groups to be infected predominantly through heterosexual relationships, similar to the findings for Group II patients in this study, while the second group was infected predominantly through homosexual relationships, as observed for the Group I patients in this study. Contact viablood was not reported, despite

being an important route of infection in Minas Gerais, resulting in over 30% of cases of HIV infection [21, 22].

The use of antiretroviral therapy (ART) was nearly 100% among patients in the Group II, demonstrating the easy access to these drugs by patients, as regulated by the Unified Health System as found recently in Brazilian northeast [23]. In comparison, none of the patients with HIV/AIDS evaluated in Vietnam were under antiretroviral therapy [24] and only approximately 15.00% of the examined patients in India were receiving ART [25]. It is widely accepted that the use of antiretroviral drugs promotes a noticeable improvement in the prognosis of HIV infection by

Table 3: Association Between the Presence of Oral Manifestation Related to HIV with TCD4+ Lymphocyte Count, TCD8 Lymphocyte Count with Age in the Group I Patients Regulated by SUS-Goiás in the Period of 2011 to 2012

Variables	Oral Manifestation							
	Absent		Present		p	95% CI		
	n	%	n	%		OR	Min	Max
CD4								
< 200 cells/mm ³	-	-	-	-				
200 a 349 cells/mm ³	20	27	2	11,1				
350 a 500 cells/mm ³	22	29,7	9	50				
> 500 cells/mm ³	32	43,2	7	38,9				
Total	74	100	18	100	0,580	1,21	(0,62-	2,34)
CD8								
<500 cells/mm ³	2	2,7	0	0				
500 a 749 cells/mm ³	11	14,9	3	16,7				
750 A 1000 cells/mm ³	20	27	6	33,3				
>1000 cells/mm ³	41	55,4	9	50				
Total	74	100	18	100	0,933	0,97	(0,52-	1,83)
Age								
18 to 39 yearsold	54	73	8	44,4				
40 to 49years old	17	23	6	33,3				
≥50 yearsold	3	4,1	4	22,2				
Total	74	100	18	100	0,008	2,82	(1,31-	6,05)

OR Odds Ratio CI Confidence Interval p significance level.

Table 4: Association between the presence of Oral Manifestations Related to HIV with Regular Dental Treatment, Use of Prostheses and Caries Presence in the Group I Patients Regulated by SUS-Goiás in the Period of 2011 to 2012

Variables	Oral Manifestation							
	Absent		Present		p	95% CI		
	n	%	n	%		OR	Min	Max
Regular Dental Treatment								
No	39	52,7	13	72,2				
Yes	35	47,3	5	27,8				
Total	74	100	18	100	0,141	0,43	(0,14-	1,32)
Use of Prostheses								
No	63	85,1	11	61,1				
Yes	11	14,9	7	38,9				
Total	74	100	18	100	0,027	3,6	(1,2-	11,4)
Caries								
No	47	63,5	10	55,6				
Yes	27	36,5	8	44,4				
Total	74	100	18	100	0,534	1,39	(0,49-	3,95)

OR Odds Ratio CI Confidence Interval p significance level.

Table 5: Association between the presence of Oral Manifestation Related to HIV with Regular Dental Treatment, Use of Prostheses and Caries Presence in the Group I Patients Regulated by SUS-Goiás in the Period of 2011 to 2012

Variables	Oral Manifestation							
	Absent		Present		p	95% CI		
	n	%	n	%		OR	Min	Max
Regular Dental Treatment								
No	49	67,1	28	77,8				
Yes	24	32,9	8	22,2				
Total	73	100	36	100	0,253	0,58	(0,23-	1,47)
Use of Dental Prostheses								
No	53	72,6	23	63,9				
Yes	20	27,4	13	36,1				
Total	73	100	36	100	0,353	1,5	(0,64-	3,51)
Caries								
No	48	65,8	16	44,4				
Yes	25	34,2	20	55,6				
Total	73	100	36	100	0,035	2,4	(1,1-	5,4)

OR Odds Ratio p significance level CI Confidence Interval.

modifying AIDS-associated pathology, including manifestations in the oral cavity. Thus, oral manifestations start appearing casually presentation [26]. However, a broad use of highly potent antiretroviral wasn't enough to eradicate oral opportunistic infections in this study.

The general and individual prevalence of oral manifestations in HIV infection detected was lower than reported in 154 HIV+ patients in Cuba, with a prevalence of 40.90% for any oral lesion [27] and in a study of 76 HIV+ patients evaluated in Aracaju (Brazil), with a prevalence of 57.89% [28]. Moreover, in another Brazilian study conducted in Recife (Brazil), 33.50% of the HIV+ patients examined presented with one or more oral manifestations associated with HIV [29]. In Iran, Bakshae (2014) [22] found 95.9% of participants with at least one oral manifestation.

Patients with advanced stages of HIV infection had less competent immune systems, which are less effective at reacting to opportunistic agents; this observation may explain why the prevalence of HIV related oral manifestations in the Group II was almost twice as high as the Group I.

HIV-related periodontitis can be explained by a number of factors, such as the marked immunodeficiency in HIV/AIDS patients, a reduced lymphocyte response against pathogenic agents and the presence of bacterial pathogens on the gums.

This condition facilitates the production of pro-inflammatory cytokines that increase the chance of periodontal damage. These alterations appear early in the course of HIV infection [30]. The findings about periodontitis are consistent with those from studies with Nigerian (14.00%) [4] and Indian (85.00%) patients with HIV/AIDS who had periodontitis [30]. The occurrence of necrotizing ulcerative gingivitis in both groups was consistent with an evaluation of Nigerian patients [4]. The linear gingival erythema (LGE) affected almost twice as many patients in Group II compared with Group I, but had not been observed in India [30]. LGE aetiology seems to involve an invasion by *Candida* species of the gingival tissue [31]. These results were consistent with studies in Brazil, where periodontal disease associated with HIV was registered in 11.00% and 9.00% of the investigated patients [32], showing a clear reduction in the occurrence of these lesions once they started treatment with nonnucleoside inhibitors analogs of the reverse transcriptase [33]. It's likely that the disseminated use of antibiotics, antifungal and antiretroviral drugs used in treatment of HIV-infection has an important impact on the development of these lesions [34]. On the other hand, classic periodontal disease, unrelated to HIV infection, is quite prevalent in the general population and in affected study patients in both groups.

Different forms of candidiasis affecting the oral cavity had a somewhat low prevalence in this study

compared to studies in Africa (55.10%), Brazil (59.00%) and Thailand (55.10% of the northern population and 25.00% of the southern population) [2,15,18]. Brazilian and Australian patients with HIV/AIDS presented a higher prevalence in oral candidiasis [28, 35] and erythematous candidiasis has remained the most prevalent process in the oral cavity even with the use of highly active antiretroviral therapy (HAART) [26, 36].

Combination therapies with protease inhibitors provide a reduction in the exoenzyme production by *Candida* species, especially by *Candida albicans*, reducing the ability of these fungi to become aggressive in HIV+ patients. These patients have more *C. albicans* proteolytic enzymes than HIV-negative patients. This feature facilitates the colonization of the oral cavity with these strains of fungi, increasing the prevalence of erythematous candidiasis [37]. In Nigerian patients, more than 80.00% of the subjects with candidiasis also presented with angular cheilitis. This situation can be explained by the action of HIV in patients with a poor nutritional status, as observed for patients from Lagos (Nigeria) and for patients in many poor African countries [4].

Regarding oral viral lesions, this study differed from research conducted in Nigeria in 700 patients, where a 23.00% prevalence of herpes zoster lesions was observed, although only 35 (5.00%) of the patients were HIV-positive [4]. In Brazil, only 8.30% of patients had lesions of herpes labialis [19].

Oral hairy leukoplakia (OHL) results confirmed previous findings in Barcelona [26] and suggested a reduction in the appearance of this entity that has been related to a CD4+ cell count below 300 cells/mm³, and a viral load increase and is considered a reliable marker for the progression of HIV infection [38]. In contrast, 24.77% of Thai patients had OHL [39].

The salivary gland diseases manifested primarily as xerostomia, which occurred in some individuals receiving antiretroviral therapy as seen in Australia [35]. No cases of Kaposi's sarcoma were diagnosed, confirming similar studies in Brazil and Nigeria [15,40]. Nowadays this neoplasm is considered rare [38]. These results are in contrast to an African research (Zimbabwe) where this neoplasm was diagnosed in 18.60% of the examined patients [2].

The high prevalence of caries in the study population, especially in Group II, demonstrated the

inaccessibility to regular dental treatment. A similar result was observed in Iran and Spain [22, 26]. This situation is of concern because affects the quality of life of these patients by adversely affecting their ability to eat appropriately.

Among the adverse oral effects reported by patients, those caused by antiretroviral therapy affected both groups. In the Group II, more than 90% of patients were on antiretroviral therapy. A higher prevalence of xerostomia (47.80%) was detected among Spanish patients on HAART [26]. This condition can compromise the patient's oral health by facilitating the emergence of infections (it may lead to caries lesions). Nausea, aphthous ulcers and taste disturbances occurred in a similar prevalence reported in other research study conducted in Brazil [33].

Melanotic hyperpigmentation was infrequent; however, other studies have shown contradictory results, with this lesion having a prevalence of 34.00% and 19.00% in India and Nigeria, respectively [25, 40]. The introduction of HAART changes the prevalence of oral manifestations in patients even after only one month of treatment, although the positive effect of this therapy does not seem to provide sufficient protection against xerostomia or lesions induced by the human papillomavirus (HPV) [10,41].

The absence of a significant association between the occurrence of oral manifestation related to HIV and the CD4+ lymphocyte count was opposite to that which was observed in other studies [18, 28, 32]. An African study found an association between CD4+ cell counts of <500 cells/mm³ and the emergence of oral candidiasis [42]. No significant association between oral lesions and viral load was observed in this study, which is in contrast to the results obtained from another Brazilian study [32] that found a greater number of oral lesions associated with viral loads above 10.000 copies/mm³. A significant association with the presence of oral lesions and age in the Group I differed from Brazilian research that did not find such an association [32]. The use of dental prostheses associated with the presence of HIV-related oral manifestations in Group I corroborates findings from another Brazilian study, which suggested that the rough surface of the resin used for dental prostheses accumulate microorganisms, especially *Candida albicans*. These devices therefore require careful hygiene to prevent the buildup of bacterial plaques that may function as niches for opportunistic pathogens [37]. In the Group II, the presence of caries was significantly associated with the

onset of HIV-related oral lesions, possibly due to the virulence of microorganisms involved in cariogenesis and the severe disruption of the immune system in these more advanced stages of HIV infection.

A significant difference between the HIV+ and AIDS Groups is not surprising because the immune status of patients with AIDS sharply deteriorates during this advanced stage of HIV infection.

In conclusion, Group II patients were more affected than Group I patients by oral lesions. Periodontitis was the most prevalent oral lesion. Age and prosthesis-wearing were influential factors for the development of these lesions in Group I patients, while the presence of caries influenced oral lesion development in Group II patients.

One limitation of this study was the difficulty in confirming the presumptive diagnosis by means of laboratory tests, which may lead to misdiagnosis. However, oral lesions associated with HIV are easily diagnosed during a clinical examination, which minimizes the possibility of misdiagnosis. We must consider the region-specific geographic aspects and health policies for a study population. Ideally, there should be standard methodologies for similar investigations across geographic locations, which would reduce discrepancies and residual differences in the results.

Finally, these findings would alert dentists about their role in address a preventive treatment for these patients.

ACKNOWLEDGEMENTS

The authors thank The Infectious Diseases Ambulatory team of Hospital of Clinics for collaboration and comprehension about this research.

CONFLICTS OF INTEREST

The authors declared no conflicts of interest.

REFERENCES

- [1] Ranganathan K, Hemalatha R. Oral lesions in HIV infection in developing countries: an overview. *Adv Dent Res* 2006; 19: 63-8. <http://dx.doi.org/10.1177/154407370601900113>
- [2] Chidzonga MM. HIV/AIDS orofacial lesions in 156 Zimbabwean patients at referral oral and maxillofacial surgical clinics. *Oral Dis* 2003; 9: 317-22. <http://dx.doi.org/10.1034/j.1601-0825.2003.00962.x>
- [3] Urizar JMA, Goicouría MAE, Del Valle AE. Síndrome de Inmunodeficiencia Adquirida: Manifestaciones en la cavidade bucal. *Med Oral Patol Oral Cir Bucal* 2004; 9: 148-57.
- [4] Agbelusi GA, Wright AA. Oral lesion as indicators of HIV infection among routine in Lagos, Nigeria. *Oral Dis* 2005; 11: 370-3. <http://dx.doi.org/10.1111/j.1601-0825.2005.01132.x>
- [5] Tamí-Maury I, Coulibaly YI, Cissoko SS, Dao S, Kristensen S. First report of HIV- related oral manifestations in Mali. *Pan Afr Med J* 2012; 11: 18.
- [6] Cherry-Peppers G, Daniel CO, Meeks V, Sanders CF, Reznik D. Oral Manifestations in The Era of HAART. *J Nat Med Assoc* 2003; 95(Suppl 2): 21S-32S.
- [7] Aleixo RQ, Scherma AP, Guimarães G, Cortelli JR, Cortelli SC. DMFT index and oral mucosal lesions associated with HIV infection: cross sectional study in Porto Velho, Amazonian Region - Brazil. *Braz J Infect Dis* 2010; 14: 449-56. [http://dx.doi.org/10.1016/S1413-8670\(10\)70092-1](http://dx.doi.org/10.1016/S1413-8670(10)70092-1)
- [8] Prasad HK, Bhojwani KM, Shenoy V, Prasad SC. HIV manifestations in otolaryngology. *Am J Otolaryngol* 2006; 27(3): 179-85. <http://dx.doi.org/10.1016/j.amjoto.2005.09.011>
- [9] Reisacher WR, Finn DG, Stern J, Zeifer B, Cocker R. Manifestations of AIDS in the head and neck. *South Med J* 1999; 92(7): 684-97. <http://dx.doi.org/10.1097/00007611-199907000-00007>
- [10] Liberali SA, Coates EA, Freeman AD, Logan RM, Jamieson L, Mejia G. Oral conditions and their social impact among HIV dental patients, 18 years on. *Australian Dental Assoc* 2012; 58: 18-25. <http://dx.doi.org/10.1111/adj.12031>
- [11] Fernandez-Cruz E, Desco M, Garcia Montes M, Longo N, Gonzalez B, Zabay JM. Immunological and serological markers predictive of progression to AIDS in a cohort of HIV-infected drug users. *AIDS* 1990; 4: 987-94. <http://dx.doi.org/10.1097/00002030-199010000-00007>
- [12] National Center for Infectious Disease of HIV /AIDS Castro KG, Ward JW, Slutsker L, Buehler JW, Jaffe HW, Berkelman RL. 1993 Revised Classification System for HIV Infection and Expanded Surveillance Case Definition for AIDS Among Adolescents and Adults. CDC 1993. Available in www.cdc.gov/. Access in 05/02/2012.
- [13] EC- Clearinghouse on oral problems related to HIV infection and WHO Collaborating Centre on oral manifestations of the immunodeficiency virus. Classification and diagnostic criteria for oral lesions in HIV infection. *J Oral Pathol Med* 1993; 22: 289-91. <http://dx.doi.org/10.1111/j.1600-0714.1993.tb01074.x>
- [14] Agwu E, Ihongbe JC, Tirwomwe JF, Pazos V, Tirwomwe M, Casadesus L. Appraisal of oral lesions status of HIV/AIDS patients in south western Uganda. *Braz J Oral Sci* 2008; 7: 1591-5.
- [15] Gasparin AB, Ferreira FV, Danesi CC, et al. Prevalência e fatores associados às manifestações bucais em pacientes HIV positivos atendidos em cidade sul-brasileira. *CadSaude Pública* 2009; 25: 1307-15. <http://dx.doi.org/10.1590/S0102-311X2009000600013>
- [16] Khongkuntian P, Grote M, Isaranatan W, Plyaworawong S, Reichart PA. Oral manifestations in HIV-positive adults from northern Thailand. *J Oral Pathol Med* 2001; 30: 220-3. <http://dx.doi.org/10.1034/j.1600-0714.2001.300405.x>
- [17] Bendick C, Schelfele C, Reichart PA. Oral manifestations in 102 cambodian patients with HIV infection and AIDS. *J Oral Pathol Med* 2002; 31: 1-4. <http://dx.doi.org/10.1034/j.1600-0714.2002.310101.x>
- [18] Kerdpon D, Pongsiriwet S, Pangsomboon K, et al. Oral manifestations of HIV infection in relation to clinical CD4 immunological status in northern and southern Thai patients. *Oral Dis* 2004; 10: 138-44. <http://dx.doi.org/10.1046/j.1601-0825.2003.00990.x>

- [19] Amorim JA, Souza FMB, Costa EB, Carneiro VSM, De Lucena AAG. Prevalência das doenças estomatológicas em pacientes HIV positivos. *Odontol Clin – Científ* 2009; 8: 127-31.
- [20] Vajpayee M, Kanswal P, Seth P, Wiq N. Spectrum of opportunistic infections and profile of CD4+ Counts among AIDS Patients in North India. *Infection* 2003; 31: 336-40.
- [21] Nobre V, Braga E, Rayes A, *et al.* Opportunistic infections in southeast of Brazil. *Rev Inst Med Trop São Paulo* 2003; 45: 69-4.
<http://dx.doi.org/10.1590/S0036-46652003000200003>
- [22] Bakhshae M, Sarvghad MR, Khazaeni K, Movahed R, Hoseinpour AM. HIV: An Epidemiologic study on head and neck Involvement in 50 Patients. *Iranian J Otorhinolaryngol* 2014; 26(2): 97-03.
- [23] Motta WKS, Nóbrega DRM, Santos MGC, Gomes DKC, Godoy GP, Pereira JV. Aspectos demográficos e Manifestações clínicas bucais de pacientes soropositivos para o HIV/Aids. *Rev Odontol UNESP* 2014; 43(1): 61-67.
<http://dx.doi.org/10.1590/S1807-25772014000100010>
- [24] Jensen TO, Tam VV, Mai NT, *et al.* Oral and constitutional manifestations of HIV-infected hospital patients in northern Vietnam. *Southeast Asian J Trop Med Public Health* 2005; 36: 1459-68.
- [25] Sharma G, Pai KM, Suhas S, Ramapuram JT, Doshi D, Anup N. Oral manifestations in HIV/AIDS infected patients from India. *Oral Dis* 2006; 12: 537-42.
<http://dx.doi.org/10.1111/j.1601-0825.2006.01232.x>
- [26] Jané-Salas E, Chimenos-Kustner E, López-López J, Roselló-Llabrés X, Ocaña-Rivera I. Efecto de los tratamientos antirretrovirales en las manifestaciones orales de los pacientes VIH+. *Avances en Odontología* 2006; 22: 315-26.
<http://dx.doi.org/10.4321/S0213-12852006000600003>
- [27] Carpio E, López V, Fardales V, Benítez I. Oral manifestations of HIV infection in adults patients from province of Sancti Spiritus, Cuba. *J Oral Pathol Med* 2009; 38: 126-31.
<http://dx.doi.org/10.1111/j.1600-0714.2008.00716.x>
- [28] Medeiros CF, Santos TB, Albuquerque Júnior RLC, Moura SAB. Relação entre as manifestações estomatológicas, contagem de células CD4+ e carga viral em pacientes HIV positivos. *Pesq Bras Odontoped Clin Integr* 2007; 7: 271-6.
<http://dx.doi.org/10.4034/1519.0501.2007.0073.0013>
- [29] Pinheiro A, Marcenes W, Zakrzewska JM, Robinson PG. Dental and oral lesions in HIV infected patients: a study in Brazil. *Int Dent J* 2004; 54: 131-7.
<http://dx.doi.org/10.1111/j.1875-595X.2004.tb00268.x>
- [30] Ranganathan K, Magesh KT, Kumarasamy N, Solomon S, Viswanathan R, Johnson NW. Greater severity and extent of periodontal breakdown in 136 south indian human immunodeficiency virus seropositive patients than in normal controls: A comparative study using community periodontal index of treatment needs. *Indian J Dent Res* 2007; 18: 55-9.
<http://dx.doi.org/10.4103/0970-9290.32420>
- [31] Gómez RS, da Costa JE, Loyola AM, de Araújo NS, de Araújo VC. Immunohistochemical study of linear gingival erythema from HIV-positive patients. *J Periodontol Res* 1995; 30: 355-9.
<http://dx.doi.org/10.1111/j.1600-0765.1995.tb01287.x>
- [32] Mattos SL, Santos VR, Ferreira EF. Prevalência de lesões de mucosa bucal em pacientes HIV positivos da unidade de referência especializada em doenças infecciosas e parasitárias especiais – URE- DIPE (Belém-Pará). URL: <http://www.patologiaoral.com.br/texto46.asp> (accessed in November 18, 2012).
- [33] Ortega KL, Vale DA, Magalhães MHCG. Impact of PI and NNRTI HAART – based therapy on oral lesions of Brazilian HIV-infected patients. *J Oral Pathol Med* 2009; 38: 489-94.
<http://dx.doi.org/10.1111/j.1600-0714.2009.00783.x>
- [34] Gonçalves LS, Gonçalves BML, Fontes TV. Periodontal disease in HIV-infected adults in the HAART era: Clinical, immunological, and microbiological aspects. *Arch Oral Biol* 2013; 58: 1385-96.
<http://dx.doi.org/10.1016/j.archoralbio.2013.05.002>
- [35] Freeman AD, Liberali SA, Coates EA, Logan RM. Oral health in Australian HIV patients since the advent of combination antiretroviral therapy. *Aust Dent J* 2012; 57: 470-476.
<http://dx.doi.org/10.1111/adj.12005>
- [36] Gaurav S, Keerthilatha PM, Archana N. Prevalence of Oral Manifestations and Their Association with CD4/CD8 Ratio and HIV Viral Load in South India. *Int J Dent* 2011; 1-8.
<http://dx.doi.org/10.1155/2011/964278>
- [37] Aleva NA, Birman EG, Afonso Jr. W, *et al.* Erythematous candidosis in patients with complete dentures and HIV/AIDS. *Mycoses* 2007; 50: 407-11.
<http://dx.doi.org/10.1111/j.1439-0507.2007.01390.x>
- [38] Miziara ID, Lima AS, De La Cortina RAC. Candidíase oral e leucoplasia pilosa como marcadores de progressão da infecção pelo HIV em pacientes brasileiros. *Rev Bras Otorrinolaringol* 2004; 70(3): 310-4.
<http://dx.doi.org/10.1590/S0034-72992004000300005>
- [39] Pattrapornnan P, DeRouen TA. Associations of periodontitis and oral manifestations with CD4 counts in HIV-pregnant women in Thailand. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2013; 116(3): 306-12.
<http://dx.doi.org/10.1016/j.ooolo.2013.04.016>
- [40] Adedigba MA, Ogunbodede EO, Jeboda SO, Naidoo S. Patterns of oral manifestation of HIV/AIDS among 225 Nigerian patients. *Oral Dis* 2008; 14: 341-6.
<http://dx.doi.org/10.1111/j.1601-0825.2007.01384.x>
- [41] Schmidt-Westhausen AM, Pripke F, Bergmann FJ, Reichart PA. Decline in the rate of oral opportunistic infections following introduction of highly active antiretroviral therapy. *J Oral Pathol Med* 2000; 29: 336-41.
<http://dx.doi.org/10.1034/j.1600-0714.2000.290708.x>
- [42] Adurogbangba MI, Aderinokun GA, Odaibo GN, Olaleye OD, Lawoyin TO. Orofacial lesions and CD4 counts associated with HIV/AIDS in an adult population in Oio State, Nigeria. *Oral Dis* 2004; 10: 319-26.
<http://dx.doi.org/10.1111/j.1601-0825.2004.01036.x>